CYANIDE 113

3. CHEMICAL AND PHYSICAL INFORMATION

3.1 CHEMICAL IDENTITY

Information regarding the chemical identity of cyanide is located in Table 3-1. Hydrogen cyanide is a toxic gas which enters the environment from both natural processes and human industrial activities. It may exist in polymeric forms. The cyanide compounds in which cyanide can be obtained as CN are classified as simple and complex cyanides. Some simple cyanides are soluble in water (sodium cyanide, NaCN; potassium cyanide, KCN; calcium cyanide, Ca(CN)₂), while others are sparingly soluble or almost insoluble (copper (I) cyanide, CuCN). Cyanogen chloride (CNC₁) is a highly toxic gas that is soluble in water. At alkaline pH, CNC1 hydrolyzes to CNO which has only limited toxicity. Alkaline chlorination of water containing cyanide produces cyanogen chloride. Thiocyanate (SCN) is an oxidation product of cyanide, produced in the presence of sulfur.

3.2 PHYSICAL AND CHEMICAL PROPERTIES

Information regarding the physical and chemical properties of cyanide is located in Table 3-2. Cyanides form strong complexes with many metals, particularly those of the transition series. One example of such complexation is the reaction of cyanide with iron in the formation of ferrocyanide and ferricyanide complexes. Solutions of ferrocyanides and ferricyanides can form hydrogen cyanide and cyanide ions when exposed to sunlight or ultraviolet radiation. Cyanogenic glycosides are cyanide compounds produced naturally in many plants. These glycosides produce hydrogen cyanide when hydrolyzed (Towill et al. 1978).

Hydrogen cyanide has a pK_a of 9.2 (Smith and Martell 1989); therefore, solutions of cyanide compounds in water (such as from sodium cyanide and potassium cyanide) can form hydrogen cyanide at acid and neutral pHs. Alkaline solutions with pH >12 are practical for preventing significant outgassing of hydrogen cyanide.

Hydrogen cyanide is a fire hazard and may be explosive when an excess of a strong acid is added to confined hydrogen cyanide. Solutions of some cyanide compounds are not stable and may decompose upon exposure to air or light.

Table 3-1. Chemical Identity of Cyanide and Compounds^a

Characteristic	Hydrogen cyanide	Sodium cyanide	Potassium cyanide	Calcium cyanide
Synonym(s)	Formonitrile; hydrocyanic acid; prussic acid	Cyanide of sodium; hydrocyanic acid; sodium salt	Cyanide of potassium; hydrocyanic acid, potassium salt	Calcid; calcyan; cyanide of calcium
Registered trade name(s)	Cyclone B; Cyclon ^b	Cyanogran ^c	Carswell No. 688A	Caswell No. 142 Cyanogas ^c
Chemical formula	HCN	NaCN	KCN	Ca(CN) ₂
Chemical structure	$H^+C \equiv N^-$	Na ⁺ C≡N ⁻	K+C≡N-	$^{-}N \equiv CCa^{+2}C \equiv N^{-}$
Identification numbers:				
CAS registry	74–90–8	143–33–9	151–50–8	592-01-8
NIOSH RTECS	MW6825000	VZ7530000	TS8750000	EW0700000
EPA hazardous waste	P063, D003	P106; D003	P098; D003	P021; D003
OHM/TADS	7216749	7216892	7216862	7216626
DOT/UN/NA/IMCO shipping	UN1051; IMO 6.1 UN1614; NA 1051	UN1689; IMO 6.1	UN1680; IMO 6.1	UN1575; IMO 6.1
HSDB	165	734	1245	242
NCI	No data	No data	No data	No data

Table 3-1. Chemical Identity of Cyanide and Compounds^a (continued)

Characteristic	Copper(I) cyanide	Potassium silver cyanide	Cyanogen	Cyanogen chloride
Synonym(s)	Cuprous cyanide ^c ; cupricin ^c	Potassium argento- cyanide; potassium dicyanoargentate	Carbon nitride; dicyanogen; ethanedinitrile	Chlorine cyanide; chlorocyan
Registered trade name(s)	AI3-28745	No data	No data	Caswell No. 267
Chemical formula	CuCN	KAg(CN) ₂	(CN) ₂	CNCI
Chemical structure	Cu ⁺ C≡N⁻	K ⁺ [Ag(CN) ₂] ⁻	$N \equiv C - C \equiv N$	CI-C≡N
Identification numbers:				
CAS registry	544-92-3	506-61-6	460–19–5	506–77–4
NIOSH RTECS	GL7150000	TT5775000	GT1925000	GT2275000
EPA hazardous waste	P029; D003	P099; D003; D011	P031; D003	P033; D003
OHM/TADS	No data	No data	7216656	7216658
DOT/UN/NA/IMCO shipping	UN1587; IMO 6.1	No data	UN1026; IMO 2.3	UN1589; IMO 2.3
HSDB	1438	6053	2130	917
NCI	No data	No data	No data	No data

Table 3-1. Chemical Identity of Cyanide and Compounds^a (continued)

Characteristic	Ammonium thiocyanate	
Synonym(s)	Thiocyanic acid, ammonium salt; ammonium rhodanide; ammonium sulfocyanate ^c	
Registered trade name(s)	Trans-Aid ^b	
Chemical formula	$NH_4^+S-C\equiv N^-$	
Chemical structure	NH₄SCN	
Identification numbers:	4	
CAS registry	1762-95-4	
NIOSH RTECS	XK7875000	
EPA hazardous waste	No data	
OHM/TADS	721218	
DOT/UN/NA/IMCO shipping	NA9092	
HSDB	701	
NCI	No data	

^aAll data are from HSDB 1996 unless otherwise noted.

CAS = Chemical Abstracts Service; DOT/UN/NA/IMO = Dept. of Transportation/United Nations/North America/International Maritime Dangerous Goods Code; EPA = Environmental Protection Agency; HSDB = Hazardous Substances Data Bank; NCI = National Cancer Institute; NIOSH = National Institute for Occupational Safety and Health; OHM/TADS = Oil and Hazardous Materials/Technical Assistance Data System; RTECS = Registry of Toxic Effects of Chemical Substances

^bFarm Chemicals Handbook 1983

^cMerck 1989

Table 3-2. Physical and Chemical Properties of Cyanide and Compounds

Property	Hydrogen cyanide	Sodium cyanide
Molecular weight	27.03 ^a	49.01 ^a
Color	Colorless ^h	White ^h ; colorless ^a
Physical state	Gas or liquid ^h	Solid ^h
Melting point, °C	-13.4 ^h	563.7 ^a
Boiling point, °C	25.70 ^d	1496 ^a
Density, g/cm ³	0.6884 (liquid at 20 °C) ^d ;	1.60 (for cubic form) ^d
Odor	Faint bitter almond odor ^c	Odorless when dry, emits slight odor of HCN in damp air ^h
Odor threshold:		
Water	0.17 ppm (w/v) ^b	No data
Air	$0.58 \text{ ppm } (v/v)^{b}; 0.8-4.4 \text{ ppm}^{g}$	No data
Solubility:		40 4400 L 140 00d
Water Organic solvent(s)	Miscible ^a Soluble in ethanol, ether ^a	48 g/100 mL at 10 °C ^d Slightly soluble in ethanol ^a and formamide ^d
Partition coefficients:		
Log K _{ow}	0.66 ^e ; 1.07 (calc.) ^l	0.44 ^e
Log K _{oc}	No data	No data
Vapor pressure, mm Hg	630 (at 20 °C) ^g	0.76 at 800 °C ^d
Henry's law constant	5.1x10 ⁻² atm-m ³ /mol ^f	No data
Autoignition temperature	538 ^d	No data
Flashpoint, °C	–17.8 (closed cup) ^d	No data
Flammability limits	5.6–40% ⁿ	Not combustible ⁿ
Conversion factors: mg/m ³ to ppm in air, 20 °C ppm to mg/L in water ppm to mg/kg in soluble samples	1 mg/m ³ = 0.890 ppm ppm (w/v) = mg/L = μg/mL ppm (w/w) = mg/kg = μg/g Upper, 40%; lower, 5.6% ⁹	NA ⁱ ppm (w/v) = mg/L = μg/mL ppm (w/w) = mg/kg = μg/g No data

3. CHEMICAL AND PHYSICAL INFORMATION

Table 3-2. Physical and Chemical Properties of Cyanide and Compounds (continued)

Property	Potassium cyanide	Calcium cyanide
Molecular weight	65.12 ^a	92.12 ^a
Color	White ^h ; colorless ^a	White ^a
Physical state	Solid ^h	Solid ^a
Melting point, °C	634.5 ^a	Decomposes at >350 °Ca
Boiling point, °C	No data	No data
Density, g/cm ³	1.553 (for cubic form) ^d	1.8–1.9 (commercial product) ^d
Odor	Faint bitter almond odor ^h	Faint bitter almond odor ^h
Odor threshold:		
Water	No data	No data
Air	No data	No data
Solubility: Water	71.6 g/100 mL at 25 °C ^d	Soluble in water with gradual liberation of HCN ^h
Organic solvent(s)	Slightly soluble in ethanol ^d and methanol ^h	
Partition coefficients:		
Log K _{ow}	No data	No data
Log K _{oc}	3.0 (calc.) ^m	No data
Vapor pressure, mm Hg	No data	No data
Henry's law constant	No data	No data
Autoignition temperature	No data	No data
Flashpoint	No data	No data
Flammability limits	Not combustible ⁿ	Not combustible ⁿ
Conversion factors: mg/m³ to ppm in air, 20 °C ppm to mg/L in water ppm to mg/kg in soluble samples	NA ⁱ ppm (w/v) = mg/L = μg/mL ppm (w/w) = mg/kg = μg/g	NA ⁱ ppm (w/v) = mg/L = μg/mL
Explosive limits	No data	No data

3. CHEMICAL AND PHYSICAL INFORMATION

Table 3-2. Physical and Chemical Properties of Cyanide and Compounds (continued)

Property	Potassium silver cyanide	Cyanogen
Molecular weight	199.01 ^h	52.04 ^a
Color	White ^h	Colorless ^a
Physical state	Solid ^h	Gas ^a
Melting point, °C	No data	–27.9 ^a
Boiling point, °C	No data	-20.7 ^a
Density, g/cm ³	2.36 ^a	0.9577 at21.17 °C ^a
Odor	No data	Almond-like odor ^h
Odor threshold:		
Water	No data	No data
Air	No data	230 ppm; irritating at 15 ppm ⁹
Solubility:		
Water	Soluble ^h	450 cc/100 cc (20 °C) ^a
Organic solvent(s)	Slightly soluble in ethanol ^a	Soluble in ethanol and ethyl ether ^a
Partition coefficients:		
Log K _{ow}	No data	No data
Log K _{oc}	No data	No data
Vapor pressure, mm Hg	No data	3,800 at 20 °C ^j
Henry's law constant	No data	No data
Autoignition temperature	No data	No data
Flashpoint	No data	No data
Flammability limits	No data	6.6 – 32% in air ⁿ
Conversion factors: mg/m³ to ppm in air, 20 °C ppm to mg/L in water ppm to mg/kg in soluble samples	NA^{i} ppm (w/v) = mg/L = μ g/mL ppm (w/w) = mg/kg = μ g/g	1 mg/m ³ = 0.462 ppm ppm (w/v) = mg/L = μg/mL ppm (w/w) = mg/kg = μg/g
Explosive limits	No data	No data

3. CHEMICAL AND PHYSICAL INFORMATION

Table 3-2. Physical and Chemical Properties of Cyanide and Compounds (continued)

Property	Cyanogen chloride	Copper(I) cyanide
Molecular weight	61.47 ^a	89.56 ^a
Color	Colorless ^c	White to cream-coloredh
Physical state	Gas ^c	Solid ^a
Melting point, °C	-6 ^a	473 (in N ₂) ^a
Boiling point, °C	13.8 ^h ;12.7 ^a	Decomposesa
Density, g/cm ³	1.186 ^h	2.92 ^a
Odor	Highly irritating ^h	No data
Odor threshold:		
Water	No data	No data
Air	1 ppm ^g	No data
Solubility: Water	Soluble ^h	Insoluble ^a
Organic solvent(s)	Soluble Soluble in ethanol and ethyl ether ^h	Insoluble in alcohol ^g
Partition coefficients:		
Log K _{ow}	No data	No data
Log K _{oc}	No data	No data
Vapor pressure, mm Hg	760 at 13.8 °C	No data
Henry's law constant	No data	No data
Autoignition temperature	No data	No data
Flashpoint	No data	No data
Flammability limits	Not combustible ^g	Does not readily ignite ^g
Conversion factors: mg/m ³ to ppm in air, 20 °C ppm to mg/L in water ppm to mg/kg in soluble samples	1 mg/m ³ = 2.5 ppm ppm (w/v) = mg/L = μg/mL ppm (w/w) = mg/kg = μg/g	NA ⁱ ppm (w/v) = mg/L = μg/mL ppm (w/w) = mg/kg = μg/g

Table 3-2. Physical and Chemical Properties of Cyanide and Compounds (continued)

Property	Ammonium thiocyanate
Molecular weight	76.12 ^a
Color	Colorless ^a
Physical state	Solid ^a
Melting point, °C	149.6 ^a
Boiling point, °C	170 decomposes ^a
Density, g/cm ³	1.305 ^a
Odor	Odorless ^h
Odor threshold: Water Air	No data No data
Solubility: Water	128 g/100 cc at 0 °C; v soluble in hot water ^a
Organic solvent(s)	v soluble in ethanol; soluble in acetone and methanol; insoluble in ethyl acetate and chloroform ^h
Partition coefficients:	
Log K _{ow} Log K _{oc}	No data No data
Vapor pressure, mm Hg	No data
Henry's law constant	No data
Autoignition temperature	No data
Flashpoint	No data
Flammability limits	May be combustible ^g
Conversion factors: mg/m ³ to ppm in air, 20 °C ppm to mg/L in water ppm to mg/kg in soluble samples	NA^{i} ppm (w/v) = mg/L = μ g/mL ppm (w/w) = mg/kg = μ g/g
Explosive limits	No data

^aLide 1993

^bAmoore and Hautala 1983

^cHawley 1981

^dJenks 1979

^eEPA 1984a

^fYoo et al. 1986; value at 25 °C and saturation pressure

⁹HSDB 1994

^hMerck 1989

ⁱSince these compounds do not exist in the atmosphere in the vapor phase, their concentrations are always expressed in weight by volume unit (e.g., mg/m³).

Towill et al. 1978

^kSax 1984

^IVerschueren 1983 ^mKenaga 1980 ⁿNFPA 1994

EPA = Environmental Protection Agency; HCN = hydrogen cyanide; HSDB = Hazardous Substances Data Bank; NA = not applicable

		·	